

Second, there are additional statistical challenges to these tests. DerSimonian and Laird's [8] Q statistic is specific to random effects models, while Cochran's Q statistic [7] is not presented in the context of covariates: their original derivation is only for models that do not control for confounding. This is presumably because the authors are well aware that one cannot directly compare magnitudes of odds ratios from separate analyses, a point only recently highlighted in the applied literature [11–13]. In addition, because of differences in the weighting of trials when calculating Cochran's Q for different summary statistics, these comparisons may not be particularly meaningful and may be misleading [10].

Third, in practice estimating a model with a rich specification of covariates can provide a good approximation to other models. We showed this in another article in which multivariate logistic regression could recover the adjusted risk ratio even when the data generating process is an odds ratio [14]. DerSimonian and Laird [8] also argue that although choosing the wrong measure could imply heterogeneity in treatment effects, in practice this would not tend to happen unless the rate for the control group varied widely or was close to either zero or one.

Fourth, our preferred approach to choosing a measure is to begin with the policy or research question. Even if this approach requires tolerating reduced precision because of heterogeneity, we hold with Tukey [15], "Far better an approximate answer to the right question, which is often vague, than an exact answer to the wrong question, which can always be made precise." We consider the statistics to serve the substance and not vice versa.

We thank Valkenhoef and Ades for their careful read of our article, and for drawing attention to important issues about conducting meta-analyses. Their letter is a great reminder that assumptions matter, that some statistical models are theoretically incompatible with seemingly similar models, and that getting the right statistical model is extremely important. We hope that this exchange will generate better understanding of the strengths and weaknesses of different approaches to indirect comparison meta-analyses.

Edward C. Norton, PhD

Department of Health Management and Policy

Department of Economics

University of Michigan and National Bureau of Economic Research

Ann Arbor, MI, USA

Morgen M. Miller, BS

Department of Health Management and Policy

Department of Economics, University of Michigan

Ann Arbor, MI, USA

Jason Wang, PhD

Department of Health Evidence and Policy

Mount Sinai School of Medicine

New York, NY, USA

Kasey Coyne, BA

Department of Health Evidence and Policy

Mount Sinai School of Medicine

New York, NY, USA

Lawrence C. Kleinman, MD, MPH

Departments of Health Evidence and Policy and Pediatrics

Mount Sinai School of Medicine

New York, NY, USA

1098-3015/\$36.00 – see front matter Copyright © 2013,  
International Society for Pharmacoeconomics and Outcomes  
Research (ISPOR). Published by Elsevier Inc.  
<http://dx.doi.org/10.1016/j.jval.2013.02.003>

## REFERENCES

- [1] van Valkenhoef G, Ades AE. Evidence synthesis assumes additivity on the scale of measurement: response to "rank reversal in indirect comparisons" by Norton et al. *Value Health* 2013;16:449–52.
- [2] Norton EC, Miller MM, Wang JJ, et al. Rank reversal in indirect comparisons. *Value Health* 2012;15:1137–40.
- [3] Lu G, Ades AE. Modeling between-trial variance structure in mixed treatment comparisons. *Biostatistics* 2009;10:792–805.
- [4] Ades AE. ISPOR states its position on network meta-analysis. *Value Health* 2011;14:414–6.
- [5] Lu G, Welton NJ, Higgins JPT, et al. Linear inference for mixed treatment comparison meta-analysis: a two-stage approach. *Res Synth Methods* 2011;2:43–60.
- [6] Caldwell DM, Welton NJ, Dias S, Ades AE. Selecting the best scale for measuring treatment effect in a network meta-analysis: a case study in childhood nocturnal enuresis. *Res Synth Methods* 2012;3:126–41.
- [7] Cochran WG. The combination of estimates from different experiments. *Biometrics* 1954;10:101–29.
- [8] DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177–88.
- [9] Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–58.
- [10] Deeks JJ. Issues in the selection of a summary statistic for meta-analysis of clinical trials with binary outcomes. *Stat Med* 2002;21:1575–600.
- [11] Mroz TA, Zayats YV. Arbitrarily normalized coefficients, information sets, and false reports of "biases" in binary outcome models. *Rev Econ Stat* 2008;90:406–13.
- [12] Mood C. Logistic regression: why we cannot do what we think we can do, and what we can do about it. *Eur Soc Rev* 2010;26:67–82.
- [13] Norton EC. Log odds and ends. NBER Working Paper #18252, 2012.
- [14] Kleinman LC, Norton EC. What's the risk? A simple approach for estimating adjusted risk ratios from nonlinear models including logistic regression. *Health Serv Res* 2009;44:288–302.
- [15] Tukey JW. The future of data analysis. *Ann Math Stat* 1962;33:1–67.

## ERRATUM

In *Value in Health* Volume 16, Issue 1, pages 46–56, (January/February 2013), the correct authors listing for the following article should be as follows:

### Cost-Effectiveness of Gene-Expression Profiling For Tumor-Site Origin

John Homberger, MD, MS, Irina Degtiar, BS, BA, Hialy Gutierrez, BS, Ashwini Shewade, MS, MSc, W. David Henner, PhD, MD, Shawn Becker, MD, Gauri Varadhachary, MD, Stephen Raab, MD